

## Recommended Adult Immunization Schedule, UNITED STATES, OCTOBER 2005–SEPTEMBER 2006

- 1. Tetanus and Diphtheria (Td) vaccination.** Adults with uncertain histories of a complete primary vaccination series with diphtheria and tetanus toxoid-containing vaccines should receive a primary series using combined Td toxoid. A primary series for adults is 3 doses; administer the first 2 doses at least 4 weeks apart and the third dose 6–12 months after the second. Administer 1 dose if the person received the primary series and if the last vaccination was received  $\geq 10$  years previously. Consult ACIP statement for recommendations for administering Td as prophylaxis in wound management ([www.cdc.gov/mmwr/preview/mmwrhtml/00041645.htm](http://www.cdc.gov/mmwr/preview/mmwrhtml/00041645.htm)). The American College of Physicians Task Force on Adult Immunization supports a second option for Td use in adults: a single Td booster at age 50 years for persons who have completed the full pediatric series, including the teenage/young adult booster. A newly licensed tetanus-diphtheria-acellular pertussis vaccine is available for adults. ACIP recommendations for its use will be published.
- 2. Measles, Mumps, Rubella (MMR) vaccination.** *Measles component:* adults born before 1957 can be considered immune to measles. Adults born during or after 1957 should receive  $\geq 1$  dose of MMR unless they have a medical contraindication, documentation of  $\geq 1$  dose, history of measles based on healthcare provider diagnosis, or laboratory evidence of immunity. A second dose of MMR is recommended for adults who 1) were recently exposed to measles or in an outbreak setting, 2) were previously vaccinated with killed measles vaccine, 3) were vaccinated with an unknown type of measles vaccine during 1963–1967, 4) are students in postsecondary educational institutions, 5) work in a healthcare facility, or 6) plan to travel internationally. Withhold MMR or other measles-containing vaccines from HIV-infected persons with severe immunosuppression. *Mumps component:* 1 dose of MMR vaccine should be adequate for protection for those born during or after 1957 who lack a history of mumps based on healthcare provider diagnosis or who lack laboratory evidence of immunity. *Rubella component:* administer 1 dose of MMR vaccine to women whose rubella vaccination history is unreliable or who lack laboratory evidence of immunity. For women of child-bearing age, regardless of birth year, routinely determine rubella immunity and counsel women regarding congenital rubella syndrome. Do not vaccinate women who are pregnant or might become pregnant within 4 weeks of receiving the vaccine. Women who do not have evidence of immunity should receive MMR vaccine upon completion or termination of pregnancy and before discharge from the healthcare facility.

- 3. Varicella vaccination.** Varicella vaccination is recommended for all adults without evidence of immunity to varicella. Special consideration should be given to those who 1) have close contact with persons at high risk for severe disease (healthcare workers and family contacts of immunocompromised persons) or 2) are at high risk for exposure or transmission (e.g., teachers of young children; child care employees; residents and staff members of institutional settings, including correctional institutions; college students; military personnel; adolescents and adults living in households with children; nonpregnant women of childbearing age; and international travelers). Evidence of immunity to varicella in adults includes any of the following: 1) documented age-appropriate varicella vaccination (i.e., receipt of 1 dose before age 13 years or receipt of 2 doses [administered at least 4 weeks apart] after age 13 years); 2) born in the United States before 1966; 3) history of varicella disease based on healthcare provider diagnosis or self- or parental report of typical varicella disease for non-U.S.-born persons born before 1966 and all persons born during 1966–1997 (for a patient reporting a history of an atypical, mild case, healthcare providers should seek either an epidemiologic link with a typical varicella case or evidence of laboratory confirmation, if it was performed at the time of acute disease); 4) history of herpes zoster based on healthcare provider diagnosis; or 5) laboratory evidence of immunity. Do not vaccinate women who are pregnant or might become pregnant within 4 weeks of receiving the vaccine. Assess pregnant women for evidence of varicella immunity. Women who do not have evidence of immunity should receive dose 1 of varicella vaccine upon completion or termination of pregnancy and before discharge from the healthcare facility. Dose 2 should be given 4–8 weeks after dose 1.

- 4. Influenza vaccination.** *Medical indications:* chronic disorders of the cardiovascular or pulmonary systems, including asthma; chronic metabolic diseases, including diabetes mellitus, renal dysfunction, hemoglobinopathies, or immunosuppression (including immunosuppression caused by medications or by HIV); any condition (e.g., cognitive dysfunction, spinal cord injury, seizure disorder or other neuromuscular disorder) that compromises respiratory function or the handling of respiratory secretions or that can increase the risk of aspiration; and pregnancy during the influenza season. No data exist on the risk for severe or complicated influenza disease among persons with asplenia; however, influenza is a risk factor for secondary bacterial infections that can cause severe disease among persons with asplenia. *Occupational indications:* healthcare workers and employees of long-term care and assisted living facilities. *Other indications:* residents of nursing homes and other long-term care and assisted living facilities; persons likely to transmit influenza to persons at high risk (i.e., in-home household contacts and caregivers of children birth through 23 months of age, or persons of all ages with high-risk conditions); and anyone who wishes to be vaccinated.



## Footnotes

# Recommended Adult Immunization Schedule, UNITED STATES, OCTOBER 2005—SEPTEMBER 2006

For healthy nonpregnant persons aged 5–49 years without high-risk conditions who are not contacts of severely immunocompromised persons in special care units, intranasally administered influenza vaccine (FluMist®) may be administered in lieu of inactivated vaccine.

**5. Pneumococcal polysaccharide vaccination.** *Medical indications:* chronic disorders of the pulmonary system (excluding asthma); cardiovascular diseases; diabetes mellitus; chronic liver diseases, including liver disease as a result of alcohol abuse (e.g., cirrhosis); chronic renal failure or nephrotic syndrome; functional or anatomic asplenia (e.g., sickle cell disease or splenectomy [if elective splenectomy is planned, vaccinate at least 2 weeks before surgery]); immunosuppressive conditions (e.g., congenital immunodeficiency, HIV infection [vaccinate as close to diagnosis as possible when CD4 cell counts are highest], leukemia, lymphoma, multiple myeloma, Hodgkin disease, generalized malignancy, organ or bone marrow transplantation); chemotherapy with alkylating agents, antimetabolites, or high-dose, long-term corticosteroids; and cochlear implants. *Other indications:* Alaska Natives and certain American Indian populations; residents of nursing homes and other long-term care facilities.

**6. Revaccination with pneumococcal polysaccharide vaccine.** One-time revaccination after 5 years for persons with chronic renal failure or nephrotic syndrome; functional or anatomic asplenia (e.g., sickle cell disease or splenectomy); immunosuppressive conditions (e.g., congenital immunodeficiency, HIV infection, leukemia, lymphoma, multiple myeloma, Hodgkin disease, generalized malignancy, organ or bone marrow transplantation); or chemotherapy with alkylating agents, antimetabolites, or high-dose, long-term corticosteroids. For persons aged  $\geq 65$  years, one-time revaccination if they were vaccinated  $\geq 5$  years previously and were aged  $< 65$  years at the time of primary vaccination.

**7. Hepatitis A vaccination.** *Medical indications:* persons with clotting factor disorders or chronic liver disease. *Behavioral indications:* men who have sex with men or users of illegal drugs. *Occupational indications:* persons working with hepatitis A virus (HAV)-infected primates or with HAV in a research laboratory setting. *Other indications:* persons traveling to or working in countries that have high or intermediate endemicity of hepatitis A (for list of countries, visit [www.cdc.gov/travel/diseases.htm#hepa](http://www.cdc.gov/travel/diseases.htm#hepa)) as well as any person wishing to obtain immunity. Current vaccines should be given in a 2-dose series at either 0 and 6–12 months, or 0 and 6–18 months. If the combined hepatitis A and hepatitis B vaccine is used, administer 3 doses at 0, 1, and 6 months.

**8. Hepatitis B vaccination.** *Medical indications:* hemodialysis patients (use special formulation [40  $\mu$ g/mL] or two 20- $\mu$ g/mL doses) or patients who receive clotting factor concentrates. *Occupational indications:* health-care workers and public-safety workers who have exposure to blood in the workplace; and persons in training in schools of medicine, dentistry, nursing, laboratory technology, and other allied health professions. *Behavioral indications:* injection-drug users; persons with more than one sex partner in the previous 6 months; persons with a recently acquired sexually transmitted disease (STD); and men who have sex with men. *Other indications:* household contacts and sex partners of persons with chronic hepatitis B virus (HBV) infection; clients and staff of institutions for the developmentally disabled; all clients of STD clinics; inmates of correctional facilities; or international travelers who will be in countries with high or intermediate prevalence of chronic HBV infection for  $> 6$  months (for list of countries, visit [www.cdc.gov/travel/diseases.htm#hepa](http://www.cdc.gov/travel/diseases.htm#hepa)).

**9. Meningococcal vaccination.** *Medical indications:* adults with anatomic or functional asplenia, or terminal complement component deficiencies. *Other indications:* first-year college students living in dormitories; microbiologists who are routinely exposed to isolates of *Neisseria meningitidis*; military recruits; and persons who travel to or reside in countries in which meningococcal disease is hyperendemic or epidemic (e.g., the “meningitis belt” of sub-Saharan Africa during the dry season [Dec–June]), particularly if contact with the local populations will be prolonged. Vaccination is required by the government of Saudi Arabia for all travelers to Mecca during the annual Hajj. Meningococcal conjugate vaccine is preferred for adults meeting any of the above indications who are aged  $\leq 55$  years, although meningococcal polysaccharide vaccine (MPSV4) is an acceptable alternative. Revaccination after 5 years may be indicated for adults previously vaccinated with MPSV4 who remain at high risk for infection (e.g., persons residing in areas in which disease is epidemic).

**10. Selected conditions for which *Haemophilus influenzae* type b (Hib) vaccine may be used.** *Haemophilus influenzae* type b conjugate vaccines are licensed for children aged 6 weeks–71 months. No efficacy data are available on which to base a recommendation concerning use of Hib vaccine for older children and adults with the chronic conditions associated with an increased risk for Hib disease. However, studies suggest good immunogenicity in patients who have sickle cell disease, leukemia, or HIV infection, or have had splenectomies; administering vaccine to these patients is not contraindicated.

# Recommended Adult Immunization Schedule, by Vaccine and Age Group UNITED STATES, OCTOBER 2005--SEPTEMBER 2006

Vaccine	Age group	19-49 years	50-64 years	≥ 65 years
Tetanus, diphtheria (Td) <sup>1*</sup>		1 or 2 doses	1-dose booster every 10 yrs	1 dose
Measles, mumps, rubella (MMR) <sup>2*</sup>		2 doses (0, 4-8 wks)	2 doses (0, 4-8 wks)	1 dose
Varicella <sup>3*</sup>		1 dose annually	1 dose annually	1 dose annually
Influenza <sup>4*</sup>		1-2 doses	1-2 doses	1 dose
Pneumococcal (polysaccharide) <sup>5,6</sup>		2 doses (0, 6-12 mos, or 0, 6-18 mos)	2 doses (0, 6-12 mos, or 0, 6-18 mos)	1 dose
Hepatitis A <sup>7*</sup>		3 doses (0, 1-2, 4-6 mos)	3 doses (0, 1-2, 4-6 mos)	1 or more doses
Hepatitis B <sup>8*</sup>		1 or more doses	1 or more doses	1 or more doses
Meningococcal <sup>9</sup>		1 or more doses	1 or more doses	1 or more doses

--- Vaccines below broken line are for selected populations

NOTE: These recommendations must be read along with the footnotes.  
\*Covered by the Vaccine Injury Compensation Program.

For all persons in this category who meet the age requirements and who lack evidence of immunity (e.g., lack documentation of vaccination or have no evidence of prior infection)

Recommended if some other risk factor is present (e.g., based on medical, occupational, lifestyle, or other indications)

This schedule indicates the recommended age groups and medical indications for routine administration of currently licensed vaccines for persons aged ≥ 19 years. Licensed combination vaccines may be used whenever any components of the combination are indicated and when the vaccine's other components are not contraindicated. For detailed recommendations, consult the manufacturers' package inserts and the complete statements from the ACIP ([www.cdc.gov/nip/publications/acip-list.htm](http://www.cdc.gov/nip/publications/acip-list.htm)).

Report all clinically significant postvaccination reactions to the Vaccine Adverse Event Reporting System (VAERS). Reporting forms and instructions on filing a VAERS report are available by telephone, 800-822-7967, or from the VAERS website at [www.vaers.hhs.gov](http://www.vaers.hhs.gov).

Information on how to file a Vaccine Injury Compensation Program claim is available at [www.hrsa.gov/osp/vicp](http://www.hrsa.gov/osp/vicp) or by telephone, 800-338-2382. To file a claim for vaccine injury, contact the U.S. Court of Federal Claims, 717 Madison Place, N.W., Washington D.C. 20005, telephone 202-357-6400.

Additional information about the vaccines listed above and contraindications for vaccination is also available at [www.cdc.gov/nip](http://www.cdc.gov/nip) or from the CDC-INFO Contact Center at 800-CDC-INFO (232-4636) in English and Spanish, 24 hours a day, 7 days a week.



DEPARTMENT OF HEALTH AND HUMAN SERVICES  
CENTERS FOR DISEASE CONTROL AND PREVENTION



# Recommended Adult Immunization Schedule, by Vaccine and Medical and Other Indications

## UNITED STATES, OCTOBER 2005–SEPTEMBER 2006

Vaccine ▼	Indication ►	Pregnancy	Congenital immunodeficiency; leukemia; <sup>11</sup> lymphoma; generalized malignancy; cerebrospinal fluid leaks; therapy with alkylating agents, antimetabolites, radiation, or high-dose, long-term corticosteroids	Diabetes; heart disease; chronic pulmonary disease; chronic liver disease, including chronic alcoholism	Asplenia <sup>10</sup> (including elective splenectomy and terminal component deficiencies)	Kidney failure, end-stage renal disease, recipients of hemodialysis or clotting factor concentrates	Human immunodeficiency virus (HIV) infection; <sup>12</sup>	Healthcare workers
Tetanus, diphtheria (Td) <sup>1*</sup>					1-dose booster every 10 yrs			
Measles, mumps, rubella (MMR) <sup>2*</sup>					1 or 2 doses			
Varicella <sup>3*</sup>					2 doses (0, 4–8 wks)			2 doses
Influenza <sup>4*</sup>					1 dose annually		1 dose annually	
Pneumococcal (polysaccharide) <sup>5,6</sup>		1–2 doses			1–2 doses			1–2 doses
Hepatitis A <sup>7*</sup>				2 doses (0, 6–12 mos, or 0, 6–18 mos)				
Hepatitis B <sup>8*</sup>			3 doses (0, 1–2, 4–6 mos)				3 doses (0, 1–2, 4–6 mos)	
Meningococcal <sup>9</sup>			1 dose		1 dose		1 dose	

NOTE: These recommendations must be read along with the footnotes.

\*Covered by the Vaccine Injury Compensation Program.

For all persons in this category who meet the age requirements and who lack evidence of immunity (e.g., lack documentation of vaccination or have no evidence of prior infection)

Recommended if some other risk factor is present (e.g., based on medical, occupational, lifestyle, or other indications)

Contraindicated

Approved by the Advisory Committee on Immunization Practices (ACIP), the American College of Obstetricians and Gynecologists (ACOG), and the American Academy of Family Physicians (AAFP)